

INFLUENZA PANDEMICS DURING THE 20TH CENTURY

- An influenza pandemic is a global outbreak of disease that occurs when a new influenza A virus appears or “emerges” in the human population, causes serious illness, and then spreads easily from person to person worldwide.
- Pandemics are different from seasonal outbreaks or “epidemics” of influenza. Seasonal outbreaks are caused by subtypes of influenza viruses that already circulate among people, whereas pandemic outbreaks are caused by new subtypes, by subtypes that have never circulated among people, or by subtypes that have not circulated among people for a long time.

During the 20th century, the emergence of several new influenza A virus subtypes caused three pandemics, all of which spread around the world within a year of being detected. The pandemics led to high levels of illness, death, social disruption, and economic loss.

- **1918-19**, "Spanish flu," [A (H1N1)], caused the highest number of known influenza deaths. (However, the actual influenza virus subtype was not detected in the 1918-19 pandemic). More than 500,000 people died in the United States, and up to 50 million people may have died worldwide. Many people died within the first few days after infection, and others died of secondary complications. Nearly half of those who died were young, healthy adults. Influenza A (H1N1) viruses still circulate today after being introduced again into the human population in 1977.
- **1957-58**, "Asian flu," [A (H2N2)], caused about 70,000 deaths in the United States. First identified in China in late February 1957, the Asian flu spread to the United States by June 1957.
- **1968-69**, "Hong Kong flu," [A (H3N2)], caused about 34,000 deaths in the United States. This virus was first detected in Hong Kong in early 1968 and spread to the United States later that year. Influenza A (H3N2) viruses still circulate today.

Both the 1957-58 and 1968-69 pandemics were caused by viruses containing a combination of genes from a human influenza virus and an avian influenza virus. The 1918-19 pandemic virus appears to have an avian origin.

PREPARING FOR THE NEXT PANDEMIC

Many scientists believe it is only a matter of time until the next influenza pandemic occurs. The severity of the next pandemic cannot be predicted, but modeling studies suggest that the impact of a pandemic on the United States could be substantial. In the absence of any control measures (vaccination or drugs), it has been estimated that in the United States a “medium-level” pandemic could cause 89,000 to 207,000 deaths, 314,000 and 734,000 hospitalizations, 18 to 42 million outpatient visits, and another 20 to 47 million people being sick. Between 15% and 35% of the U.S. population could be affected by an influenza pandemic, and the economic impact could range between \$71.3 and \$166.5 billion.